

Derivatives Practice – Math 210/211

Power Rule: If $f(x) = ax^n$, then $f'(x) = anx^{n-1} \cdot dx$.

Note: this rule applies to radicals (fractional exponents) and simple rational expressions (negative exponents).

The “ dx ” that trails is meant to signify the chain rule. If “ x ” (the base) happens to be a function itself, then you must apply the chain rule. Technically, you always apply the chain rule, but if the base is simply “ x ”, then the chain rule gives “1” as the derivative, and the answer is unchanged.

Example: If $f(x) = 4x^3$, then $f'(x) = 12x^2 \cdot 1$. The chain rule gives a “1” since it’s the derivative of x , but we usually just write the answer as $f'(x) = 12x^2$.

Example: If $f(x) = 3(x^2 + 2)^5$, then $f'(x) = 15(x^2 + 2)^4 \cdot (2x)$. In this case, the chain rule gives a derivative of $2x$, which trails the answer. We would normally clean up the answer and write: $f'(x) = 30x(x^2 + 2)^4$.

Exponential Rule: If $f(x) = e^x$, then $f'(x) = e^x \cdot dx$.

Example: $f(x) = e^{2x} \Rightarrow f'(x) = 2e^{2x}$.

Natural Logarithm Rule: If $f(x) = \ln x$, then $f'(x) = \frac{1}{x} dx = \frac{dx}{x}$.

Example: $f(x) = \ln(2x^3 + x + 1) \Rightarrow f'(x) = \frac{1}{2x^3 + x + 1} (6x^2 + 1) = \frac{6x^2 + 1}{2x^3 + x + 1}$.

Note how the final answer is combined into one expression for convenience.

Product Rule: If $f(x) = u \cdot v$, then $f'(x) = u \cdot v' + v \cdot u'$.

Quotient Rule: If $f(x) = \frac{u}{v}$, then $f'(x) = \frac{v \cdot u' - u \cdot v'}{v^2}$.

Practice:

- 1) $f(x) = x^2 + 3x + 1$
- 2) $f(x) = 4x^6 + 1.5x^4 + 3x^3 - x + 17$
- 3) $f(x) = 2(x^2 + x)^2$
- 4) $f(x) = \sqrt{3 + x^2}$
- 5) $f(x) = \sqrt[3]{x^2 + 1}$
- 6) $g(x) = \frac{3}{x^3}$
- 7) $g(x) = \frac{4}{(x^2 + 1)^2}$
- 8) $g(x) = \frac{5}{2x^4}$
- 9) $h(x) = e^{3x^2 + x + 2}$
- 10) $h(x) = (e^{2x} + 1)^4$
- 11) $j(x) = \ln(x + x^2 + 2x^3)$
- 12) $j(x) = \ln(e^x + 2)$
- 13) $j(x) = x \ln(x + 1)$
- 14) $k(x) = e^{3x} \ln x$
- 15) $k(x) = (\ln(x + 1) + x)^3$
- 16) $k(x) = \frac{e^{5x}}{\ln(2x)}$
- 17) $m(x) = \frac{x^2 + 1}{(x^3 + 1)^2}$
- 18) $m(x) = \frac{e^{2x} \ln x}{x^2}$
- 19) $n(x) = 25^2 + e$
- 20) $p(x) = \sqrt{\frac{x}{e^x}}$

Answers

(In no particular order!)

(Some have been simplified!)

$$\frac{-4x^4 - 6x^2 + 2x}{(x^3 + 1)^3}$$

$$8e^{2x}(e^{2x} + 1)^3$$

$$\frac{x}{x+1} + \ln(x+1)$$

$$-\frac{10}{x^5}$$

$$2x + 3$$

$$-\frac{16x}{(x^2 + 1)^3}$$

$$\frac{e^x}{e^x + 2}$$

$$(6x + 1)e^{3x^2 + x + 2}$$

$$\frac{e^{3x}}{x} + 3e^{3x} \ln x$$

$$-\frac{9}{x^4}$$

$$\frac{1 + 2x + 6x^2}{x + x^2 + 2x^3}$$

$$2\sqrt{\frac{e^x}{x} \left(\frac{1-x}{e^x} \right)}$$

$$\frac{x}{\sqrt{3 + x^2}}$$

$$8x^3 + 12x^2 + 4x$$

$$0$$

$$24x^5 + 6x^3 + 9x^2 - 1$$

$$\frac{2x}{3\sqrt[3]{3 + x^2}}$$

$$\frac{3(x+2)(\ln(x+1) + x)^2}{x+1}$$

$$\frac{5xe^{5x} \ln(2x) - e^{5x}}{x(\ln(2x))^2}$$

Any errors (real or imagined), contact surgent@asu.edu. Good luck! (Updated 1/21/06)