



ELSEVIER

3 January 2002

PHYSICS LETTERS B

Physics Letters B 524 (2002) 21–25

www.elsevier.com/locate/npe

Primordial inflation and present-day cosmological constant from extra dimensions

Carl L. Gardner¹*Department of Mathematics, Arizona State University, Tempe, AZ 85287, USA*

Received 31 May 2001; received in revised form 30 August 2001; accepted 30 August 2001

Editor: J. Frieman

Abstract

A semiclassical gravitation model is outlined which makes use of the Casimir energy density of vacuum fluctuations in extra compactified dimensions to produce the present-day cosmological constant as $\rho_\Lambda \sim M^8/M_P^4$, where M_P is the Planck scale and M is the weak interaction scale. The model is based on $(4 + D)$ -dimensional gravity, with $D = 2$ extra dimensions with radius $b(t)$ curled up at the ADD length scale $b_0 = M_P/M^2 \sim 0.1$ mm. Vacuum fluctuations in the compactified space perturb b_0 very slightly, generating a small present-day cosmological constant.

The radius of the compactified dimensions is predicted to be $b_0 \approx k^{1/4} 0.09$ mm (or equivalently $M \approx 2.4$ TeV/ $k^{1/8}$), where the Casimir energy density is k/b^4 .

Primordial inflation of our three-dimensional space occurs as in the cosmology of the ADD model as the inflaton $b(t)$, which initially is on the order of $1/M \sim 10^{-17}$ cm, rolls down its potential to b_0 . © 2002 Published by Elsevier Science B.V.

PACS: 98.80.Cq; 04.50.+h; 11.25.Mj

Keywords: Cosmological constant; Inflationary cosmology; Extra dimensions

1. Introduction

Supernova data indicate that the energy density ρ_Λ in a present-day cosmological constant is on the order of $0.7\rho_c$, where the current critical density $\rho_c \approx (2.5 \times 10^{-3} \text{ eV})^4$. It is intriguing that $\rho_\Lambda \sim b_0^{-4}$ where $b_0 \sim 0.1$ mm—just the length scale for compactified extra dimensions predicted by Arkani-Hamed–Dimopoulos–Dvali (ADD) type theories [1] with two extra spatial dimensions.

It is possible that this dark energy derives from vacuum fluctuations in extra compactified dimen-

sions. We outline here a semiclassical gravitation model which makes use of this mechanism to produce the present-day cosmological constant. The model is based on $(4 + D)$ -dimensional gravity, with $D = 2$ extra dimensions with radius $b(t)$ curled up at the ADD length scale b_0 , where the subscript “0” denotes present-day values.

The ADD model can be realized [2] in type I ten-dimensional string theory, with standard model fields naturally restricted to a 3-brane [3], while gravitons propagate in the full higher-dimensional space. For $D = 2$, two of the six compactified dimensions are curled up with radius $\sim b_0$, while the remaining four are curled up with radius $\sim 1/M_I$, with the type I string scale $M_I \sim 1$ TeV. In this picture, the ADD model is formulated within a consistent quantum theory of gravity.

E-mail address: gardner@asu.edu (C.L. Gardner).

¹ Research supported in part by the National Science Foundation under grant DMS-9706792.

In addition, if supersymmetry is broken only on the 3-brane, then the bulk cosmological constant vanishes (see, e.g., Ref. [4]). A single fine tuning of parameters in the potential for b can then cancel the brane tension, setting the usual four-dimensional cosmological constant to zero.

Semiclassical $(4 + D)$ -dimensional gravitation—with a potential for the scale b of the extra compactified dimensions—rapidly becomes a good approximation to the string theory for energies below M_I [5]. In the semiclassical gravitation model, we will assume a potential for $b(t)$ which stabilizes $b(t_0)$ at $b_0 = M_P/M^2$ and which vanishes² at b_0 in the absence of the Casimir effect, where the (reduced) Planck scale $M_P = 2.4 \times 10^{18}$ GeV and the weak interaction scale $M \sim 1$ TeV. Vacuum fluctuations in the compactified space will then perturb $b(t_0)$ very slightly away from b_0 , generating a small present-day cosmological constant in our three-dimensional world. This mechanism differs from previous cosmological models incorporating the Casimir effect from vacuum fluctuations in extra compactified dimensions (see, e.g., Ref. [6]), in which the Casimir energy density in our three-dimensional world is cancelled by a bulk cosmological constant.

Primordial inflation of our three-dimensional space will occur in the model as the inflaton $b(t)$, which initially is on the order of $1/M \sim 10^{-17}$ cm, rolls down its potential to b_0 [7,8]. Many e-folds of inflation of our 3-space can occur for sufficiently flat potentials.

We will take the spacetime metric to be $R^1 \times S^3 \times T^2$ symmetric³ where S^3 is a 3-sphere and T^2 is a 2-torus:

$$g_{MN} = \text{diag}\{1, -a^2(t)\tilde{g}_{ij}, -b^2(t)\tilde{g}_{mn}\}, \quad (1)$$

where M, N run from 0 to 5; i, j run from 1 to 3; and m, n run from 4 to 5. \tilde{g}_{ij} is the metric of a unit 3-sphere and \tilde{g}_{mn} is the metric of a unit 2-torus, with $a(t)$ the radius of physical 3-space and $b(t)$ the radius of the compactified space.

² In other words, we assume that the 3-brane tension is exactly cancelled in the stabilization potential at $b = b_0$.

³ Our treatment through Eq. (11) parallels that of Kolb and Turner [9].

The nonzero components of the $(4 + D)$ -dimensional Ricci tensor are

$$\begin{aligned} R_{00} &= -\left(3\frac{\ddot{a}}{a} + D\frac{\ddot{b}}{b}\right), \\ R_{ij} &= -\left(\frac{\ddot{a}}{a} + 2\frac{\dot{a}^2}{a^2} + D\frac{\dot{a}\dot{b}}{ab} + \frac{2}{a^2}\right)g_{ij}, \\ R_{mn} &= -\left(\frac{\ddot{b}}{b} + (D-1)\frac{\dot{b}^2}{b^2} + 3\frac{\dot{a}\dot{b}}{ab}\right)g_{mn}. \end{aligned} \quad (2)$$

The generalized Einstein equations are

$$R_{MN} = 8\pi\bar{G}\left(T_{MN} - \frac{T^P{}_P}{D+2}g_{MN}\right), \quad (3)$$

where $8\pi\bar{G} = 8\pi G\mathcal{V}_0 = \mathcal{V}_0/M_P^2 = \tilde{\Omega}_D/M^{D+2}$ is the $(4 + D)$ -dimensional gravitational constant, $\mathcal{V}_0 = \tilde{\Omega}_2 b_0^2$ is the volume of the compactified dimensions today, $\tilde{\Omega}_D$ denotes the volume of the unit D -torus, and T_{MN} is the energy–momentum tensor. The gravitational coupling $8\pi G = 1/(b_0^2 M^4)$ is weak in the ADD picture because b_0 is much greater than the $(4 + D)$ -dimensional Planck length $1/M$.

The nonzero components of the energy–momentum tensor are given by

$$\begin{aligned} T_{00} &= \rho, \\ T_{ij} &= -p_a g_{ij}, \\ T_{mn} &= -p_b g_{mn}. \end{aligned} \quad (4)$$

Thus $T^P{}_P = \rho - 3p_a - Dp_b$. Expressed in terms of the radii a and b , the energy density ρ , and the pressures p_a and p_b , the Einstein equations become

$$3\frac{\ddot{a}}{a} + D\frac{\ddot{b}}{b} = -\frac{8\pi\bar{G}}{D+2}[(D+1)\rho + 3p_a + Dp_b], \quad (5)$$

$$\begin{aligned} \frac{\ddot{a}}{a} + 2\frac{\dot{a}^2}{a^2} + D\frac{\dot{a}\dot{b}}{ab} + \frac{2}{a^2} \\ = \frac{8\pi\bar{G}}{D+2}[\rho + (D-1)p_a - Dp_b], \end{aligned} \quad (6)$$

$$\frac{\ddot{b}}{b} + (D-1)\frac{\dot{b}^2}{b^2} + 3\frac{\dot{a}\dot{b}}{ab} = \frac{8\pi\bar{G}}{D+2}[\rho - 3p_a + 2p_b]. \quad (7)$$

After a few e-folds of primordial inflation of our physical 3-space, the curvature term $2/a^2$ on the left-hand side of Eq. (6) will be negligible, and we will henceforth set this term to zero.

We will be looking for solutions (neglecting matter) in which physical 3-space is inflating at the present epoch during which $b(t)$ is fixed at b_0 , or in the primordial epoch just after the quantum birth of the universe during which $b(t)$ is inflating to its present value. For an inflating 3-space (without matter), $p_a = -\rho$ and the Einstein equations become

$$3\frac{\ddot{a}}{a} + D\frac{\ddot{b}}{b} = \frac{8\pi\bar{G}}{D+2}[-(D-2)\rho - Dp_b], \quad (8)$$

$$\frac{\ddot{a}}{a} + 2\frac{\dot{a}^2}{a^2} + D\frac{\dot{a}\dot{b}}{ab} = \frac{8\pi\bar{G}}{D+2}[-(D-2)\rho - Dp_b], \quad (9)$$

$$\frac{\ddot{b}}{b} + (D-1)\frac{\dot{b}^2}{b^2} + 3\frac{\dot{a}\dot{b}}{ab} = \frac{8\pi\bar{G}}{D+2}[4\rho + 2p_b]. \quad (10)$$

The energy density and pressures on the right-hand sides of Eqs. (8)–(10) are derivable from the internal energy $U = U(a, b)$:

$$\rho = \frac{U}{\mathcal{V}}, \quad p_a = -\frac{a\partial U/\partial a}{3\mathcal{V}}, \quad p_b = -\frac{b\partial U/\partial b}{D\mathcal{V}}, \quad (11)$$

where $\mathcal{V} = \Omega_3 a^3 \tilde{\Omega}_2 b^2$ is the volume of $(3+D)$ -space and Ω_3 denotes the volume of the unit 3-sphere.

We will consider a potential $V(b)$ for the radius $b(t)$ in the internal energy

$$U(a, b) = \Omega_3 a^3 M^4 V(b) \quad (12)$$

(at zero temperature) which will produce sufficient primordial inflation to solve the horizon, flatness, homogeneity, isotropy, and monopole problems, and which will stabilize b at $b_0 = M_P/M^2 \sim 0.1$ mm, with a vanishing cosmological constant. Note that if p_a is to equal $-\rho$, then U must be proportional to a^3 , and that $V(b)$ is dimensionless.

The potential $V(b)$ will generate a potential $B(b)$ with the right-hand side of the Einstein equation (10) equal to $-B'(b)/b$. If $B(b)$ is sufficiently flat near $b \sim 1/M$, then many e-folds of inflation will occur in our physical 3-space as $b(t)$ rolls from $1/M$ to b_0 .

Quantum fields will be periodic in the compactified space, producing a Casimir effect [6] in the compactified space and in our three-dimensional world. Adding a Casimir (C) term to the internal energy

$$U_C(a, b) = \Omega_3 a^3 \left(\frac{k}{b^4} + M^4 V(b) \right) \quad (13)$$

from vacuum fluctuations in the compactified space will perturb $b(t_0)$ very slightly away from b_0 and

generate a residual present-day cosmological constant $\rho_\Lambda = k/b_0^4$. The sign and magnitude⁴ of the constant k depend on the particle content and structure of the underlying quantum gravity theory. The magnitude of k may be expected to be roughly in the range 10^{-7} – 10^{-3} based on the analysis of Candelas and Weinberg [6], who calculated the one-loop Casimir contribution from massless scalar and spin- $\frac{1}{2}$ particles in $(4+D)$ -dimensional gravitation with an odd number of extra dimensions D curled up near the Planck length. In their work, k is positive for a single massless real scalar field for odd dimensions $3 \leq D \leq 19$, but may be positive or negative. For our model to produce a positive present-day cosmological constant, we will need $k > 0$.

2. Primordial inflation

In this section, we briefly review the cosmological results for primordial inflation of Refs. [7,8] for the ADD model with internal energy U , and check that the Casimir terms in the Einstein equations when U is replaced by U_C do not qualitatively change the primordial cosmological picture.

The Einstein equations with the internal energy given by U in Eq. (12) take the form

$$3\frac{\ddot{a}}{a} + 2\frac{\ddot{b}}{b} = 3\dot{H} + 3H^2 + 2\dot{H}_b + 2H_b^2 = \frac{V'(b)}{4b}, \quad (14)$$

$$\frac{\ddot{a}}{a} + 2\frac{\dot{a}^2}{a^2} + 2\frac{\dot{a}\dot{b}}{ab} = \dot{H} + 3H^2 + 2HH_b = \frac{V'(b)}{4b}, \quad (15)$$

$$\begin{aligned} \frac{\ddot{b}}{b} + \frac{\dot{b}^2}{b^2} + 3\frac{\dot{a}\dot{b}}{ab} &= \dot{H}_b + 2H_b^2 + 3HH_b \\ &= \frac{V(b)}{b^2} - \frac{V'(b)}{4b} \equiv -\frac{B'(b)}{b}, \end{aligned} \quad (16)$$

where the Hubble parameters $H \equiv \dot{a}/a$ and $H_b \equiv \dot{b}/b$. For a vanishing present-day cosmological constant, $V'(b_0) = 0$ from Eq. (15). Eq. (16) then implies $V(b_0) = 0$ to stabilize $b(t_0)$ at b_0 .

To summarize the successful phenomenology of Ref. [8]: the ADD model can produce sufficient

⁴ A logarithmic dependence $\ln(M^2 b^2)$ can be absorbed into the definition of k without changing the conclusions below.

inflation ($\gg 70$ e-folds) to solve the cosmological problems for a class of potentials $V(b)$ which satisfy

$$H^{-1} \sim H_b^{-1} \geq \frac{1}{M} \quad (17)$$

at the beginning of inflation at the quantum birth of the universe when $a \sim b \sim 1/M$, and

$$H \gg H_b, \quad \dot{H}_b \ll H^2 \quad (18)$$

during the initial stages of inflation. The correct magnitude and approximate scale invariance of density perturbations $\delta\rho/\rho = 2 \times 10^{-5}$ are created if at an intermediate stage of inflation when $b(t) \sim 10^{3/2}/M \ll b_0$, $H_b \approx H/100$. There may be a period of contraction (similar to the vacuum Kasner solutions) of our physical 3-space, but for $D = 2$, the amount of contraction of $a(t)$ is at most 7 e-folds, so the contraction phase does not invalidate the solution of the flatness problem.

Replacing U by U_C in Eq. (13) introduces Casimir terms into the Einstein equations:

$$3\dot{H} + 3H^2 + 2\dot{H}_b + 2H_b^2 = -\frac{k}{M^4 b^6} + \frac{V'(b)}{4b}, \quad (19)$$

$$\dot{H} + 3H^2 + 2HH_b = -\frac{k}{M^4 b^6} + \frac{V'(b)}{4b}, \quad (20)$$

$$\begin{aligned} \dot{H}_b + 2H_b^2 + 3HH_b &= \frac{2k}{M^4 b^6} + \frac{V(b)}{b^2} - \frac{V'(b)}{4b} \\ &\equiv -\frac{B_C'(b)}{b}. \end{aligned} \quad (21)$$

The Casimir terms do not qualitatively change the primordial inflationary period of the ADD model, since initially

$$\frac{k}{M^4 b^6} \approx kM^2 \ll M^2 \sim H^2 \sim H_b^2 \quad (22)$$

and in the intermediate stage of inflation

$$\frac{k}{M^4 b^6} \approx 10^{-9} kM^2 \ll 10^{-11} M^2 \sim H_b^2 \sim 10^{-4} H^2 \quad (23)$$

for $k \lesssim 10^{-3}$, using the estimates in Ref. [8].

3. Present-day cosmological constant

In the present epoch, the internal dimensions have a fixed radius $b(t_0) \gg 1/M$ and $H_b = 0$. Without

the Casimir terms, the static solution for $b(t_0)$ requires $V(b_0) = 0 = V'(b_0)$. In our model, vacuum fluctuations in the compactified space perturb b_0 very slightly to \tilde{b}_0 , producing a small cosmological constant in our three-dimensional world. We assume that the potential $V(b)$ is independent of the Casimir effect, so that $V(b_0)$ and $V'(b_0)$ still equal zero.

The Einstein equations with Casimir contributions for an inflating 3-space now take the form

$$3H_0^2 = -\frac{k}{M^4 \tilde{b}_0^6} + \frac{V'(\tilde{b}_0)}{4\tilde{b}_0}, \quad (24)$$

$$0 = \frac{2k}{M^4 \tilde{b}_0^6} + \frac{V(\tilde{b}_0)}{\tilde{b}_0^2} - \frac{V'(\tilde{b}_0)}{4\tilde{b}_0}. \quad (25)$$

Setting $\tilde{b}_0 = (1 + \delta)b_0$ and solving Eq. (25) to order $\delta \sim M^4/M_P^4$ yields

$$\frac{\delta}{4} V''(b_0) + O(\delta^2) = \frac{2k}{M^4 b_0^6}, \quad (26)$$

or

$$\tilde{b}_0 \approx \left(1 + \frac{8k}{M^4 b_0^6 V''(b_0)}\right) b_0 = \left(1 + O\left(\frac{kM^4}{M_P^4}\right)\right) b_0, \quad (27)$$

where $V''(b_0) \sim 1/b_0^2 = M^4/M_P^2$. Eq. (24) then predicts a present-day cosmological term

$$\begin{aligned} 3H_0^2 &= \frac{\delta}{4} V''(b_0) - \frac{k}{M^4 b_0^6} + O(\delta^2) \\ &= \frac{k}{M^4 b_0^6} + O(\delta^2) \end{aligned} \quad (28)$$

or, in other words,

$$H_0^2 = \frac{8\pi G}{3} \rho_\Lambda, \quad \rho_\Lambda = \frac{k}{b_0^4} = \frac{kM^8}{M_P^4}. \quad (29)$$

This cosmological term will $\approx 0.7\rho_c$ if $b_0 \approx k^{1/4} 0.09$ mm, or equivalently if $M \approx 2.4$ TeV/ $k^{1/8}$.

Note that the Casimir effect has caused the stabilized radius b_0 to *increase* slightly, yielding a positive present-day cosmological constant.

The canonically normalized ‘‘radion’’ field $\varphi(t) = 2M^2 b(t)$. The mass squared of the radion field is

$$m_\varphi^2 = M^4 \frac{d^2 V}{d\varphi^2} \Big|_{\varphi_0} \sim \frac{M^4}{M_P^2} \quad (30)$$

which must be positive at $\varphi_0 = 2M^2 b_0 = 2M_P$ to have a linearly stable b_0 solution [5].

The stability properties of $B_C(b)$ in Eq. (21) are the same as of $B(b)$ in Eq. (16): the respective solutions with $b(t_0) = b_0$ and \tilde{b}_0 are linearly stable if the radion mass squared is positive, since the radion mass squared including the Casimir contribution

$$m_{\varphi,C}^2 = m_\varphi^2 + \frac{5kM^8}{M_P^6} \sim \frac{M^4}{M_P^2} \left(1 + \frac{5kM^4}{M_P^4} \right) \quad (31)$$

is positive if m_φ^2 is, and are globally stable if the respective potentials $B(b)$ and $B_C(b)$ are, for example, concave upward (the simplest case), since

$$B_C(b) = B(b) + \frac{k}{2M^4 b^4} + \text{const} \quad (32)$$

is concave upward if B is.

If the number of extra dimensions D is allowed to be greater than two, the Einstein equations (24) and (25) for an inflating 3-space with static $b(t)$ change to

$$3H_0^2 = -\frac{k}{M^{D+2}\tilde{b}_0^{D+4}} - \frac{D-2}{D+2} \frac{V(\tilde{b}_0)}{M^{D-2}\tilde{b}_0^D} + \frac{V'(\tilde{b}_0)}{(D+2)M^{D-2}\tilde{b}_0^{D-1}}, \quad (33)$$

$$0 = \frac{4k}{DM^{D+2}\tilde{b}_0^{D+4}} + \frac{4}{D+2} \frac{V(\tilde{b}_0)}{M^{D-2}\tilde{b}_0^D} - \frac{2}{D(D+2)} \frac{V'(\tilde{b}_0)}{M^{D-2}\tilde{b}_0^{D-1}} \quad (34)$$

but the result for the present-day cosmological constant has the same form

$$\rho_\Lambda = \frac{k}{b_0^4} = \frac{kM^{4+8/D}}{M_P^{8/D}}, \quad (35)$$

where now b_0 satisfies $b_0^D M^{D+2} = M_P^2$. Thus ρ_Λ has the right parametric dependence M^8/M_P^4 only for $D = 2$.

4. Conclusion

The cosmological picture presented here joins smoothly onto the primordial inflation and big-bang

cosmological pictures: the quantum birth of the universe begins with a and $b \sim 1/M$. Many ($\gg 70$) e-folds of primordial inflation occur as the inflaton $b(t)$ rolls down its potential to \tilde{b}_0 . $b(t)$ then undergoes damped oscillations about \tilde{b}_0 , heating the universe up to a temperature T above the temperature for big-bang nucleosynthesis (BBN) and creating essentially all the matter and energy we see today. (See Refs. [7] and [10] for two differing views on the maximum value of T , above which the evolution of the universe in ADD-type theories cannot be described by the radiation-dominated Friedmann–Robertson–Walker model.) At this point, the universe evolves according to the standard big-bang picture, expanding and cooling, with a fixed small cosmological constant $\rho_\Lambda = k/b_0^4 \approx (2.3 \times 10^{-3} \text{ eV})^4$.

This dark energy density is much less than the BBN energy density $\sim (1 \text{ MeV})^4$ and plays a role in the evolution of the universe only recently, long after the equality of energy density $\sim (1 \text{ eV})^4$ in matter and radiation. The radius $b(t)$ of the compactified space has not changed since well before BBN.

Finally we note that if the stabilization potential $V(b)$ vanishes at its global minimum, the resolution of the cosmic coincidences of Ref. [11] is naturally realized in the Casimir effect since parametrically $\rho_\Lambda \sim M^8/M_P^4$.

References

- [1] N. Arkani-Hamed, S. Dimopoulos, G. Dvali, Phys. Lett. B 429 (1998) 263, hep-ph/9803315.
- [2] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, G. Dvali, Phys. Lett. B 436 (1998) 257, hep-ph/9804398.
- [3] J. Polchinski, Phys. Rev. Lett. 75 (1995) 4724, hep-th/9510017.
- [4] N. Arkani-Hamed, L. Hall, D. Smith, N. Weiner, Phys. Rev. D 62 (2000) 105002, hep-ph/9912453.
- [5] N. Arkani-Hamed, S. Dimopoulos, J. March-Russell, Phys. Rev. D 63 (2001) 064020, hep-th/9809124.
- [6] P. Candelas, S. Weinberg, Nucl. Phys. B 237 (1984) 397.
- [7] N. Arkani-Hamed, S. Dimopoulos, G. Dvali, Phys. Rev. D 59 (1999) 086004, hep-ph/9807344.
- [8] N. Arkani-Hamed, S. Dimopoulos, N. Kaloper, J. March-Russell, Nucl. Phys. B 567 (2000) 189, hep-ph/9903224.
- [9] E.W. Kolb, M.S. Turner, The Early Universe, Addison-Wesley, Reading, MA, 1990.
- [10] N. Kaloper, J. March-Russell, G.D. Starkman, M. Trodden, Phys. Rev. Lett. 85 (2000) 928, hep-ph/0002001.
- [11] N. Arkani-Hamed, L.J. Hall, C. Kolda, H. Murayama, Phys. Rev. Lett. 85 (2000) 4434, astro-ph/0005111.